

Phonon and plasmon excitation in inelastic electron tunneling spectroscopy of graphite

L. Vitali, M. A. Schneider, and K. Kern

Max-Planck-Institut für Festkörperforschung, Heisenbergstr.1, D-70569 Stuttgart, Germany

L. Wirtz and A. Rubio

*Department of Material Physics, University of the Basque Country,
Centro Mixto CSIC-UPV, and Donostia International Physics Center (DIPC),
Po. Manuel de Lardizabal 4, E-20018 San Sebastián, Spain*

(Dated: February 2, 2008)

The inelastic electron tunneling spectrum (IETS) of highly oriented pyrolytic graphite (HOPG) has been measured with scanning tunneling spectroscopy (STS) at 6K. The observed spectral features are in very good agreement with the vibrational density of states (vDOS) of graphite calculated from first principles. We discuss the enhancement of certain phonon modes by phonon-assisted tunneling in STS based on the restrictions imposed by the electronic structure of graphite. We also demonstrate for the first time the local excitation of surface-plasmons in IETS which are detected at an energy of 40 meV.

PACS numbers: 68.35.Ja, 68.37.Ef, 63.20.-e

Initially applied to measure inelastic tunneling currents across metal-insulator-adsorbate-metal junctions¹, Inelastic Electron Tunneling Spectroscopy (IETS) has in the recent past been revived and its concept extended to Scanning Tunneling Microscopy (STM-IETS)². Superimposed on the elastic tunneling current, the inelastic tunneling process opens up alternative tunneling channels increasing the total conductance at the onset of its excitation. Very recently, elegant experiments (e.g.^{2,3,4,5}) have been performed in detecting local inelastic electron energy-loss spectra of individual molecule adsorbed on metallic substrates. The observed features in the tunneling spectra are found to correspond to the vibrational modes of the adsorbed species. Vibrations of single molecules as large as C₆₀ have been detected⁶. Despite the increasing number of publications of STM-IETS experiments on single molecules adsorbed on metals, to our knowledge, only two studies were reported on collective vibrational excitations of surfaces^{7,8}. In both these studies graphite was used as a substrate and several spectral features remained unexplained or were attributed to vibrational excitation in the tungsten tip. However, these tip effects have not been detected in many other STM-IETS experiments. In order to resolve the pure inelastic response of a graphite surface, we have performed STM-IETS experiments on a highly oriented pyrolytic graphite crystal (HOPG) at cryogenic temperature. In our experiments the whole vibrational spectrum of HOPG from the rigid layer shear mode at ≈ 5 meV to the optical modes at ≈ 200 meV is clearly observed without being perturbed by inelastic effects from the tip. The phonon modes of graphite can be clearly identified by the direct comparison of experimental data with first principles density function theory (DFT) calculations. We report for the first time that also surface-plasmon losses can also be observed directly with STM-IETS.

Freshly cleaved HOPG has been introduced into the

vacuum system and shortly annealed at 500K. The experiments were performed at 6K with a home built UHV-STIM using a tungsten tip. The inelastic tunneling spectrum of HOPG was measured over a topographically clean and flat terrace with a lock-in technique. In this technique a sinusoidal signal with amplitude ΔV is superimposed to the applied sample bias. In IETS the second harmonic component of the modulated signal, which is proportional to the second derivative vs. bias of the tunneling current, d^2I/dV^2 , is recorded. In this way the inelastic loss features in the conductance are transformed into peaks at the onset of the loss modes for positive sample bias and into minima for negative sample bias, respectively. A typical example for HOPG is shown in fig.1.




FIG. 1: Inelastic electron tunneling spectrum of HOPG measured by STM. The phonon features (arrows) in the d^2I/dV^2 are symmetrically located with respect to the zero bias position.

We observe phonon features that are symmetrically located with respect to zero bias as expected for the inelas-

tic tunneling process. Possibly due to a particular tip-end configuration the phonon on the negative bias side are more pronounced. In Fig. 2a we present a high-resolution spectrum of the d^2I/dV^2 signal of HOPG taken at negative bias. The data was obtained with a bias modulation of $\Delta V = 10$ meV RMS at a frequency of 2.7kHz and inverted to render the phonon excitations as peaks to facilitate the comparison to theory. Since the electron density of states of HOPG is flat in the energy range close to the Fermi level, the features in the spectrum in Fig. 1 are exclusively due to inelastic processes.




FIG. 2: a) Inelastic electron tunneling spectrum of HOPG measured by STM. Inelastic excitations show up as peaks in the d^2I/dV^2 signal. b) vibrational density of states calculated by DFT-LDA. c) Phonon dispersion relation of graphite. The phonon branches are specified as following: out of plane acoustic (ZA), acoustic shear (SH), longitudinal acoustic (LA), out of plane optical (ZO), optical shear (SH*), and longitudinal optical (LO). The inset shows the first Brillouin zone of graphite with the high symmetry points. A detailed description of the observed phonon modes is given in the text. Furthermore, a plasmon excitations is identified at 40 meV.

Along with the measured inelastic spectrum, we present in Fig. 2b and c the calculated vibrational density of states (vDOS) and the phonon dispersion relation of graphite along the A- Γ -M-K directions. Graphite is a layered structure with strong carbon bonds in the plane

Phonon branch	Symmetry point	STM-IETS (meV)	Other experiments (meV)	DFT-LDA (meV)
ZA	Γ	16	16 ²⁰	15
ZA	M	58	57 ¹²	59
ZO	M	81	81 ¹²	78
SH	M	100 (weak)	100 ¹²	77
ZO	Γ	111 (dip)	108 ¹³	109
LA	K	137	134 ¹⁹	124
SH*	M	180	172 ¹²	177
LO	Γ	200	198-205 ¹²	197-202

TABLE I: Observed features in the STM-IET spectrum of HOPG in comparison with vibrational modes observed in experiment (Neutron scattering²⁰, HREELS^{12,13} and double-resonant Raman scattering¹⁹) and with first principles DFT-LDA calculations.

and weak Van der Waals-like forces between the planes. As a result of this anisotropy, the phonon spectrum of graphite covers a very wide range. We have performed first principles calculations using density-functional theory (DFT) in the local density approximation (LDA)⁹. Aim of this calculation is to resolve some of the deviations in the vDOS calculated by the different recent force-constant parametrizations for graphite^{12,13,14,15} and to have a solid foundation for the assignment of the measured peaks. The first principles calculation has been shown^{16,17,18} to yield very close agreement with phonon dispersion relations measured by high resolution electron energy loss spectroscopy (HREELS) along the Γ -M direction¹² and along the Γ -K direction¹³. Only for the acoustic shear-mode (SH) it was observed¹⁸ that at the M-point, the HREELS data of Ref.12 yields an energy which is higher by 25 meV than the *ab-initio* value. A similar underestimation of the shear-mode frequency might occur as well at the K-point where a mode with energy 134 meV was observed by double-resonance Raman scattering¹⁹. Our calculated vDOS is also at variance with the HREELS data at M¹² but is in very good agreement with the vDOS measured by neutron scattering on a powdered sample of graphite²¹ where a separate SH(M)-feature was not observed. However, since the SH K-point mode contributes only weakly to the calculated DOS the comparison to the experimental vDOS in Ref. 21 cannot answer the question whether the mode comes out too soft in the calculations.

The observed vibrational spectral features, their comparison with values obtained with other experimental techniques and their assignment to phonon modes at the various points of the surface Brillouin zone (SBZ) are summarized in Table 1. Most features coincide with peaks in the calculated vDOS which arise from the flat dispersion relation around the high-symmetry points. In particular, the higher peaks due to phonon modes around the M-point are clearly mapped. The first peak at 16 meV is in good agreement with the calculated energy of the out of plane acoustic (ZA) mode at Γ where two

neighboring planes are oscillating out-of-phase. While in the vDOS and in neutron scattering²¹ this peak is only weak, in the IETS experiment it is strongly pronounced which can be explained by the direction of oscillation which is orthogonal to the surface plane. The weak shoulder at 6 meV corresponds to the low frequency horizontal mode at Γ where two neighboring planes are oscillating out-of-phase parallel to each other. The prominent feature at 40 meV is not due to a vibrational feature but is assigned to a plasmon excitation (see below). In Ref.7, a phonon feature at 39 meV was observed which was associated with a double-phonon excitation in the surface (16 meV) and in the tungsten tip (23 meV). However, in our experiment the tip-phonon peak is clearly not present and all the observed features can therefore be assigned to surface excitation. The vibrational feature observed at 58 meV can be assigned to the ZA at the M-point. The optical out-of-plane (ZO) mode and the shear acoustic mode SH (also called transverse acoustic mode) display a saddle in the dispersion relation around the M-point adding up to a very high peak in the vDOS at 78 meV. This explains the peak at 81 meV in the measurement. However, as noted above, HREELS data¹² suggests the energy of the SH mode at M to be about 100 meV. This gives rise to the alternative explanation that the peak at 81 meV is only due to the ZO mode while the shoulder at 100 meV may be a signal of the SH mode at the M-point. The ZO mode at Γ which gives rise to a peak at 108 meV in the theoretical vDOS seems to be missing in the spectrum. Instead, the spectrum reveals actually a dip ($dI^2/dV^2 < 0$) at this energy which will be discussed below. At 124 and 150 meV two low-intensity phonon modes, the SH, and the longitudinal acoustic (LA) modes at the K-point are predicted, respectively. Experimentally we have observed a large asymmetric inelastic feature at about 137 meV. This disagreement between experiment and theory might point to an underestimation of the SH mode also at the K point. If, as could be the case for the SH mode at M¹⁸, the real energy is higher by 10 to 20 meV, the peak at 137 meV can be assigned to the SH mode at K which corresponds to the above mentioned observation of a K-point phonon at 134 meV¹⁹ by double resonance Raman spectroscopy. In the highest energy range of the spectrum two loss-features protrude the background signal at 180 and 200 meV, respectively. The first is related to the optical shear mode (SH*, also called transverse optical mode) around the M-point. The neighboring feature at 200 meV corresponds to the longitudinal optical (LO) branch which displays a strong overbending in the direction Γ -M and Γ -K giving rise to a peak in the vDOS.

In general, the measured IET spectrum follows very closely the vibrational density of states with two notable exceptions which will be discussed in the following. These exceptions are related to the question if all phonons can be excited or if the geometry of the tunneling system leads to a restriction to phonons in a certain region of the Brillouin zone, as it is the case in phonon-

assisted electronic tunneling through narrow p - n junctions of semiconductors. In e.g. Ge, the valence-band maximum is centered at Γ while the conduction-band minimum is located at the L point. This allows only vibrational modes at L to be excited by a phonon-assisted tunneling process²². The electronic structure of graphite is also quite unique being a semimetal with a band crossing only at the K-point of the SBZ. Therefore the elastic current is carried by electrons that tunnel with a relative large $k_{\parallel} = K$ ($1.7 \text{ \AA}^{-1}$). In an STM geometry the expansion of the initial state, i.e., the tip wave function in terms of surface plane-waves²³, leads to a parallel momentum distribution centered at $k_{\parallel} = 0$. Consequently the electrons of the elastic current have a low transmission probability. Electrons with any other $k_{\parallel}$ can only tunnel by exchange of finite crystal momentum with a phonon. The phonon-assisted tunneling is relevant to all the dominant features observed besides the Γ phonons at 16 and 200 meV. An enhanced tunneling contribution should especially occur for electrons with initial parallel momentum $k_{\parallel} = 0$ that tunnel to the Fermi level at the K-point by exciting (or absorbing) a K-point phonon. This explains why the peak at 137 meV which is probably due to a K-point phonon excitation shows up so dominantly in the spectrum even though the vDOS of states only displays weak features for K-point phonons.

The second deviation of the measured spectrum from the vDOS of graphite is the dip at 111 meV where the vDOS displays a peak due to the ZO phonon at Γ . The theory for inelastic tunneling spectroscopy^{24,25} distinguishes between an inelastic contribution (real phonon emission or absorption) and an elastic contribution (virtual phonon emission with subsequent reabsorption or phonon absorption with subsequent re-emission). In some cases, the elastic contribution leads to a decrease of the tunneling conductance (dI/dV) due to back-scattering of the electron²⁴. This effect has been measured recently as dips in the dI^2/dV^2 curve of O_2 on a silver surface²⁶. We believe that the dip at 111 meV is due to a similar elastic contribution at the onset of vDOS singularity corresponding to the ZO(Γ) phonon branch. In the light of the above discussion of phonon-assisted tunneling on graphite, it is also not surprising that this effect is observed for phonons at Γ . The only matching pair for a virtual phonon scattering process are electrons initially at $k_{\parallel} = K$ and phonons at Γ . All other combinations are forbidden due to the unavailability of intermediate electronic states. Furthermore, the steady reduction of $-d^2I/dV^2$ from the ZO (M) peak at 81 meV, in contrast to the vDOS, shows that symmetry of the phonon displacement (the ZO branch) should play an important role in the suppression of the IET current²⁷. We note that one of the degenerate modes of the ZO branch at Γ is the only Γ phonon mode that has A_{2u} symmetry.

In addition to the predicted phonon modes, the measured spectrum contains an energy loss feature at 40 meV, which cannot be explained by phonon excitations. Other than the previous features, this loss feature is

of purely electronic origin and corresponds to an out-of-plane oscillation of the electrons. A tunneling electron traversing a vacuum-metal interface has a significant probability of interacting with collective electron oscillations. The overlapping of the π and π^* band at the K point of the SBZ yields a low density of free carriers (i.e. electrons and holes) near the Fermi level and allows low-energy plasmon excitation. A strong temperature dependence of the excitation energy of this plasmon mode was observed in HREELS²⁸. The excitation energy shifts from 63 meV to 45 meV when going from room temperature to 150 K, respectively. This effect has been attributed to the low density of states at the Fermi level in conjunction with the thermal excitations of carriers. Thermal excitations from the valence into the conduction band can radically alter the electron population at the Fermi level and yields the strong dependence of surface plasmon energy and line width on temperature. The temperature dependence of this excitation energy was also calculated in Ref. 28. Within the random phase approximation the plasmon energy excitation was predicted to be lowered to 36 meV at 0K in rather good agreement with our experimental value. The low-energy plasmon was also observed in transmission electron energy loss spectroscopy, which also detected a second plasmon loss, assigned to an excitation polarized parallel to the layer at 128 meV²⁹. The shoulder observed around

125 meV might be assigned to this plasmon excitation.

In conclusion, the IET spectrum of HOPG has been measured by low temperature scanning tunneling spectroscopy and is compared to an ab-initio calculation of the vibrational density of states in graphite. The observed vibrational loss features can be identified with peaks in the vDOS which are due to the flat dispersion of the phonon branches around the high-symmetry points of the Brillouin zone. Additionally, a prominent plasmon feature at 40meV has been identified in the inelastic tunneling current measured with STM. In the past, the local capability of STM-IETS has been proven to be a valuable tool to access the vibrational properties of adsorbates at surfaces. In this work we have shown that STM-IETS resolves the vibrational and collective electronic excitations of graphite with high accuracy. This is an important step for the application of the method to confined geometries and individual carbon nanostructures.

Acknowledgments

L.W. and A.R. thank support from the EC research training network COMELCAN (HPRN-CT-2000-00128), Spanish MCyT(MAT2001-0946) and UPV/EHU (9/UPV 00206.215-13639/2001).

-
- ¹ E.L.Wolf, Tunneling Spectroscopy, ed. by P.K.Hansma, New York Plenum, (1982).
 - ² B.C.Stipe, M.A.Rezaei and W.Ho, Science **280**, 1732 (1998).
 - ³ Y.Kim, T.Komeda and M.Kawai, Phys. Rev. Lett. **89**, 126104 (2002).
 - ⁴ H. J. Lee and W. Ho, Science **286**, 1719 (1999).
 - ⁵ J. I. Pascual et al., Nature **423**, 525 (2003).
 - ⁶ J.I.Pascual, J.Gomez-Herrero, D.Sanchez-Portal and H.P.Rust, J.Chem.Phys. **117**, 9531 (2002).
 - ⁷ D.P.E.Smith, G.Binnig and C.F.Quate, Appl.Phys.Lett. **49**, 1641 (1986).
 - ⁸ W.Olejniczak, Z.Klusek and M.Bieniecki, Appl.Phys.A **66**, S191 (1998).
 - ⁹ The calculations have been performed with the code ABINIT¹⁰. The force constants are calculated within linear-response theory¹¹. The wave-functions are expanded in plane waves with an energy cutoff at 80 Ry. Core electrons are described by Trouiller-Martins pseudopotentials. The 1st Brillouin zone is sampled by a 16x16x6 Monkhorst-Pack grid. The phonon frequencies are converged to within 4 cm⁻¹ with respect to variation of the energy cutoff and variation of k-point sampling. The dynamical-matrix, which is the Fourier-transform of the real-space force constants, is calculated on a 14x14x6 Monkhorst-Pack grid in the reciprocal space of the phonon-wave vector q . From this, the dynamical matrix at any q is obtained by interpolation.
 - ¹⁰ The ABINIT code is a common project of the Université Catholique de Louvain, Corning Incorporated and other contributors (URL <http://www.abinit.org>).
 - ¹¹ S. Baroni et al., Rev. Mod. Phys. **73**, 515, (2001); X. Gonze, Phys. Rev. B **55**, 10337 (1997).
 - ¹² C.Oshima et al., Solid State Comm. 65.1601.1988; T. Aizawa, R. Souda, S. Otani, Y. Ishizawa and C. Oshima, Phys. Rev. B **42**, 11469 (1990).
 - ¹³ S.Siebert, R.Pues, K.-H.Rieder and A.M.Shikin, Phys.Rev. B **55**, 7927 (1997).
 - ¹⁴ R.Al-Jishi and G.Dresselhaus, Phys.Rev. B **26**, 4514 (1982).
 - ¹⁵ A.Grüneis et al., Phys.Rev. B **65**, 155405 (2002).
 - ¹⁶ G. Kresse, J. Furthmüller and J. Hafner, Europhys. Lett. **32**, 729 (1995).
 - ¹⁷ P. Pavone et al., Physica B **219-220**, 439 (1996).
 - ¹⁸ O.Dubay and G.Kresse, Phys. Rev. B **67**, 035401 (2003).
 - ¹⁹ Y.Kawashima and G.Katagiri, Phys. Rev. B **52**, 10053 (1995).
 - ²⁰ R. Nicklow, N. Wakabayashi, and H. G. Smith, Phys. Rev. B **5**, 4951 (1972).
 - ²¹ S. Rols et al., Phys. Rev. Lett. **85**, 5222 (2000).
 - ²² R. T. Payne, Phys. Rev. **139**, A570 (1965).
 - ²³ J.Tersoff and D.R.Hamann, Phys.Rev.B **31**, 805 (1985).
 - ²⁴ B. N. J. Persson and A. Baratoff, Phys. Rev. Lett. **59**, 339 (1987).
 - ²⁵ N. Lorente and M. Persson, Phys. Rev. Lett. **85**, 2997 (2000).
 - ²⁶ J. R. Hahn, H. J. Lee, and W. Ho, Phys. Rev. Lett. **85**, 1914 (2000).
 - ²⁷ N. Lorente, M. Persson, L.J. Lauhon, W. Ho, Phys. Rev. Lett. **86**, 2593 (2001)

- ²⁸ E.T.Jensen, R.E.Palmer, W.Allison and J.F.Annett, (1971).
Phys.Rev.Lett. **66**, 492 (1991).
- ²⁹ J.Geiger, H.Katterwe and B.Schröder, Z.Physik **241**, 45